



An analytic mapping property of the Dirichlet-to-Neumann operator in Helmholtz boundary problems

Karamehmedovi, Mirza

Publication date:
2016

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Karamehmedovi, M. (2016). *An analytic mapping property of the Dirichlet-to-Neumann operator in Helmholtz boundary problems*. Abstract from 27th Nordic Congress of Mathematicians, Stockholm, Sweden.
http://www.mittag-leffler.se/sites/default/files/Applied%20Mathematics_Mirza%20Karamehmedovic.pdf

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Applied Mathematics

Location: room ??

Abstracts

An analytic mapping property of the
Dirichlet-to-Neumann operator in
Helmholtz boundary problems

Mirza Karamehmedović

Department of Applied Mathematics and Computer Science,
Technical University of Denmark

The analytic version of microlocal analysis shows that if the boundary and the Dirichlet datum of a Helmholtz boundary value problem are real-analytic, then so is the corresponding Neumann datum. However, the domain of analytic continuation of the Neumann datum is, in general, unknown. We shall here relate, in terms of explicit estimates, the domains of analytic continuation of Dirichlet and Neumann boundary data for Helmholtz problems in two or more independent variables, and in neighbourhoods of planar pieces of the boundary. For this purpose, we shall characterise a special subspace of the standard pseudodifferential operators with real-analytic symbols, to which the Dirichlet-to-Neumann operator belongs. The result can be applied in the estimation of the domain of analytic continuation of solutions across planar pieces of the boundary.